

tional immune system *in vivo*. Given that the  $\alpha$  chain alone is able to restore antigen responsiveness, it is not clear why there is a preference for expression CD8 $\alpha\beta$  heterodimers rather than CD8 $\alpha\alpha$  homodimers on mature T cells. It is possible that the CD8 $\beta$  chain has a separate role during the development of  $\alpha\beta$  TCR bearing cells in the thymus. This would correlate with the observation that intra-epithelial lymphocytes, which are mainly  $\gamma\delta$ TCR bearing cells<sup>13</sup>, predominantly express CD8 $\alpha$  homodimers on their surface<sup>14</sup>.

The expression of CD4 or CD8 accessory molecules by T cells correlates strongly with whether T cells are restricted by class II or class I MHC molecules<sup>15</sup>. Studies in transgenic mice have shown that T cells expressing class I restricted TCR  $\alpha$  and  $\beta$  transgenes favour the co-expression of CD8 molecules<sup>16,17</sup>. Although this bias may be generated by co-selection of receptor and accessory molecules binding the same class I MHC structure, it is also possible that selection occurs based on preferential interactions between particular TCR and accessory molecule transmembrane or intra-cellular domains. We examined this possibility by constructing hybrid cDNA constructs between CD4 and CD8 genes which encode the extra-cellular portion of one together with either the cytoplasmic domain, or the transmembrane plus cytoplasmic domain of the other. It was previously shown that expression of CD4 molecules by DC27.10 is insufficient to restore the response to K<sup>b</sup> even when stimulator cells express both class II and class I molecules<sup>8</sup>. If there is an interaction of the anti-K<sup>b</sup> TCR with CD8 in the intra-cellular domains then the hybrids expressing CD4 externally and CD8 internally should be capable of responding to antigen.

Transfectants expressing all combinations of hybrid CD4/CD8 molecules were isolated and stained with appropriate monoclonal antibodies (Fig. 3). All of the hybrid constructs were expressed stably on the cell surface. Immunoprecipitation of these molecules (data not shown) revealed that all constructs expressing CD4 externally were monomers, in contrast to constructs expressing CD8 $\alpha$  externally, which were expressed as disulphide bonded homodimers. This indicates that interactions between CD8 external domains dictate that this molecule is expressed normally as a dimer. Transfectants which gave similar responses to anti-clonotypic antibodies were assayed for their ability to respond to antigen (Table 1). As K<sup>b</sup> transfected L cells do not express class II MHC molecules, hybrid constructs expressing CD4 external domains were assayed on H-2K<sup>b</sup> spleen cells. Despite the presence of the CD4 ligand on the stimulator cells, expression of these hybrid molecules was insufficient to restore antigen response. By contrast, transfectants expressing hybrid molecules containing CD8 $\alpha$  external domains, whether in combination with CD4 cytoplasmic domains or CD4 transmembrane and cytoplasmic domains, functioned extremely efficiently. The latter constructs were at least as effective as native CD8 $\alpha$  homodimers in conferring responsiveness to antigen. Thus, there does not appear to be preferential association of transmembrane or intracytoplasmic domains of CD8 with a class I specific cell receptor.

These data demonstrate an incremental restoration of full antigen responsiveness conferred by transfected CD8 molecules. The first step is the progress from complete absence to limited responsiveness conferred by high level expression of CD8 extracellular domains in combination with sufficient expression of TCRs. This property can be ascribed to increased avidity stabilizing the interaction between the T cell and the antigen presenting cell. It is interesting to note that in contrast to the experiments reported by Gay *et al.*<sup>18</sup>, our experimental system has an absolute requirement for both the TCR and accessory molecules to recognize class I MHC. The Désiré-1 TCR has low affinity for antigen<sup>19</sup> and may only be stabilized by CD8 external domains binding to the same class I molecules as the TCR. This requirement may not be so obvious for higher affinity TCRs.

The second increment by which CD8 influences responsiveness relates to the presence of the cytoplasmic domain and

correlates with the ability to associate with p56<sup>lck</sup>. This association has been demonstrated for both CD4 and CD8 molecules and can explain why CD4 intracellular domains are efficient substitutes for CD8 intracellular domains. These results are in contrast to hybrid CD8 molecules with class I cytoplasmic domains which are only as efficient as CD8 $\alpha'$  chains in restoring responsiveness (B. Malissen, personal communication). □

Received 5 June; accepted 25 September 1989.

- Doyle C. & Strominger, J. L. *Nature* **330**, 256 (1987).
- Norment, A. M., Salter, R. D., Parham, P., Engelhard, V. H. & Littman, D. R. *Nature* **336**, 79–81 (1988).
- Emmrich, F., Strittmatter, U. & Eichmann, K. *Proc. natn. Acad. Sci. U.S.A.* **84**, 8298–8302 (1986).
- Ledbetter, J. A. *et al. Eur. J. Immunol.* **18**, 525–532 (1988).
- Jönsson, J.-I., Boyce, N. W. & Eichmann, K. *Eur. J. Immunol.* **19**, 253–260 (1989).
- Walker, I. D. *et al. Immunol. Rev.* **82**, 47–77 (1984).
- Zamoyska, R. & Parnes, J. R. *EMBO J.* **7**, 2359–2367 (1988).
- Gabert, J. *et al. Cell* **50**, 545–554 (1987).
- Rosenstein, Y., Ratnofsky, S., Burakoff, S. J. & Herrmann, S. J. *J. exp. Med.* **169**, 149–160 (1989).
- Rudd, C. E., Trevelyan, J. M., Dasgupta, J. D., Wong, L. L. & Schlossman, S. F. *Proc. natn. Acad. Sci. U.S.A.* **85**, 5190–5196 (1988).
- Veillette, A., Bookman, M. A., Horak, E. & Bolan, J. B. *Cell* **55**, 301–308 (1988).
- Veillette, A., Bookman, M. A., Horak, E., Samelson, L. E. & Bolan, J. B. *Nature* **338**, 257–259 (1989).
- Goodman, T. & LeFrancis, L. *Nature* **33**, 855–858 (1988).
- Parrott, D. M. V. *et al. Ann. N.Y. Acad. Sci.* **409**, 307–320 (1983).
- Swain, S. L. *Immunol. Rev.* **74**, 129–142 (1983).
- Kisielow, P., Teh, H. S., Blüthmann, H. & von Boehmer, H. *Nature* **335**, 730–733 (1988).
- Sha, W. C. *et al. Nature* **336**, 73–76 (1988).
- Gay, D. *et al. Nature* **328**, 626–629 (1987).
- Hua, C., Boyer, C., Guimenezanes, A., Alpert, F. & Schmitt-Verhulst, A. M. *J. Immunol.* **136**, 1927–1936 (1986).
- Oi, V. T., Morrison, S. L., Herzenberg, L. A. & Berg, P. *Proc. natn. Acad. Sci. U.S.A.* **80**, 825–829 (1983).
- Gunning, P., Leavitt, J., Muscat, G., Ng, S.-Y. & Keddes, L. *Proc. natn. Acad. Sci. U.S.A.* **84**, 4831–4835 (1987).
- Zamoyska, R., Vollmer, A. C., Sizer, K. C., Liaw, C. W. & Parnes, J. R. *Cell* **43**, 153–163 (1985).
- Ledbetter, J. A. & Herzenberg, L. A. *Immunol. Rev.* **47**, 63–90 (1979).

ACKNOWLEDGEMENTS. We thank Hans Stauss for K<sup>b</sup> transfected L cells and discussion, Dominique Blanc, Jean Gabert and Bernard Malissen for the DC27.10 cell line and discussing their unpublished data, and Paul Travers, Lucia Casabo, Sarah Bell and Owen Williams for reading of the manuscript. S.D.G. was a recipient of a NCI fellowship, P.V.H. is supported by a postdoctoral fellowship from the Deutsche Forschungsgemeinschaft, A.V. is supported by the National Cancer Institute of Canada. J.R.P. holds an Established Investigator Award from the American Heart Association and is supported by grants from the NIH.

## Genetic imprinting suggested by maternal heterodisomy in non-deletion Prader-Willi syndrome

Robert D. Nicholls\*†‡, Joan H. M. Knoll†, Merlin G. Butler§, Susan Karam|| & Marc Lalonde\*†¶

\* Howard Hughes Medical Institute, and † Genetics Division and Mental Retardation Center, The Children's Hospital, Department of Pediatrics, Harvard Medical School, 300 Longwood Avenue, Boston, Massachusetts 02115, USA

§ Division of Genetics, Department of Pediatrics, Vanderbilt University School of Medicine, Nashville, Tennessee 37232, USA

|| Michigan State University Medical School, East Lansing, Michigan 48824, USA

**PRADER-WILLI syndrome (PWS) is the most common form of dysmorphic genetic obesity associated with mental retardation<sup>1,2</sup>. About 60% of cases have a cytological deletion of chromosome 15q11q13 (refs 2, 3). These deletions occur *de novo* exclusively on the paternal chromosome<sup>4,5</sup>. By contrast, Angelman syndrome (AS) is a very different clinical disorder and is also associated with deletions of region 15q11q13 (refs 6–8), indistinguishable from those in PWS<sup>6,8</sup> except that they occur *de novo* on the maternal chromosome<sup>6</sup>. The parental origin of the affected chromosomes 15 in these disorders could, therefore, be a contributory factor in determining their clinical phenotypes. We have now used**

‡ Present address: Department of Neuroscience, University of Florida College of Medicine, Gainesville, Florida 32610, USA.

¶ To whom correspondence should be addressed.


cloned DNA markers specific for the 15q11q13 subregion<sup>5,9,10</sup> to determine the parental origin of chromosome 15 in PWS individuals not having cytogenetic deletions; these individuals account for almost all of the remaining 40% of PWS cases. Probands in two families displayed maternal uniparental disomy for chromosome 15q11q13. This is the first demonstration that maternal heterodisomy—the presence of two different chromosome 15s derived from the mother—can be associated with a human genetic disease. The absence of a paternal contribution of genes in region 15q11q13, as found in PWS deletion cases<sup>4,5</sup>, rather than a mutation in a specific gene(s) in this region may result in expression of the clinical phenotype. Thus, we conclude that a gene or genes in region 15q11q13 must be inherited from each parent for normal human development.

Given the unusual parental origins of deletion chromosomes associated with PWS and AS, we attempted to determine the parental origins of chromosomes 15 in non-deletion cases of PWS. For each family, we performed restriction fragment length polymorphism (RFLP) analyses with nine RFLPs at seven loci specific for proximal chromosome 15q (refs 5, 11). The first family that we studied has a proband with a phenotype typical of PWS (refs 1, 2), that is, hypotonia and growth delay in infancy, short stature, hyperphagia with consequent obesity, small hands and feet, hypogonadism and mild mental retardation. This family also displays inheritance of a balanced robertsonian

translocation, t(13; 15) (Fig. 1a). Association of this translocation with the phenotype of PWS is highly unlikely because unaffected maternal relatives carry the same balanced translocation (Fig. 1a), and the translocation breakpoint is at the centromere of chromosome 15, some distance from the critical region at 15q11q13. In this family the PWS proband (PWS1) had inherited two maternal alleles but no paternal allele (Fig. 1b) specific for probe 3-21, which maps to the region absent in PWS patients with deletions<sup>5,9,10</sup>. The 8.2-kilobase (kb) *TaqI* fragment detected by probe 3-21 (Fig. 1b) is shared by the three individuals (PWS1, his mother M1, and half-sister) carrying the t(13; 15) translocation and is, therefore, a marker for this chromosome. The 8.9-kb *TaqI* fragment which hybridized to probe 3-21 is present in both PWS1 and his mother and is not present in the other individuals of the family, clearly indicating the maternal derivation of the non-translocated chromosome 15 in PWS1, at least for region 15q11q13 surrounding the probe 3-21 locus. We confirmed and extended this result by using CMW-1, a multiallelic DNA marker<sup>11</sup> that maps distal to the region of deletions associated with PWS (unpublished result). The segregation pattern again showed two maternal alleles but no paternal allele for the proband (Fig. 1c). We excluded the possibility of non-paternity by analysis with a cloned fragment (3'HVR) from the 3' hypervariable region of the  $\alpha$ -globin locus on chromosome 16 (ref. 12, Fig. 1d).

FIG. 1 Maternal uniparental disomy in a PWS patient with a balanced Robertsonian translocation. a, Normal chromosome 15 and translocated homologue from patient PWS1 (left) and his mother M1 (right). Other unaffected maternal relatives, including the half-sister of PWS1 carry the same balanced chromosome 13q15q robertsonian translocation. b, Maternal origin of the two chromosomes 15 in patient PWS1. An 8.2-kb *TaqI* allele of probe 3-21 (D15S10) is shared by PWS1 (lane 3), his mother M1 (lane 2) and half-sister (lane 1), whereas an 8.9-kb band is shared by PWS1 and M1. PWS1 does not inherit a 9.0-kb paternal allele (F1, lane 4). Mixing experiments (lane 5, PWS1 and F1; lane 6, M1 and F1) show the identity of the three alleles in this new RFLP system. c, Presence of two maternal alleles only for CMW-1 (D15S24) in PWS1. PWS1 (lane 2) shares two *TaqI* restriction fragments, of 2.1 and 2.2 kb, with M1 (lane 3) but has neither of the 1.5- or 1.8-kb paternal alleles (F1, lane 1). The half-sister (lane 4) displays the maternal 2.1-kb and a 2.15-kb allele. The band at 1.7 kb is constant, C. d, Exclusion of non-paternity for PWS1. *PvuII*-digested DNA was hybridized to the  $\alpha$ -globin 3'HVR marker (D16S85), one of the most highly polymorphic markers in the human genome<sup>12</sup>. PWS1 (lane 2) shows mendelian inheritance of one paternal 3.9-kb allele (lane 1) and one maternal 4.3-kb allele (lane 3). The half-sister inherits her mother's 5.3-kb allele (result not shown).

METHODS. DNA was isolated from blood and/or cell lines from each individual as described previously<sup>5</sup>. Hybridizations were carried out as previously described<sup>5</sup>. High-resolution chromosome analyses from peripheral blood lymphocytes<sup>29</sup> were performed by Giemsa-trypsin banding<sup>30</sup>.



The proband (PWS2) of the second family also has classical PWS (refs 13, 14). Cytogenetically, PWS2 has two intact chromosome 15s (Fig. 2a). We demonstrated the lack of paternal alleles for two loci by haplotype analysis (Fig. 2b, c). With probe 34, the father is homozygous for 6.5 kb alleles, whereas the mother is homozygous for 6.3 kb alleles (Fig. 2b). For the proband, we found only a 6.3-kb band (Fig. 2b), consistent with the maternal origin of both cytogenetically normal chromosomes 15. Likewise, only maternal alleles specific for probe IR39d had been inherited (Fig. 2c). By dosage analyses we demonstrated that PWS2 has two copies of each of these maternally derived alleles (Fig. 2d and results not shown), indicating that loci detected

by probes 34 and IR39d had not been deleted. These two loci are flanked by loci detected by probe IR4-3R (results not shown) and probe IR10-1, respectively, (Fig. 2b) that are heterozygous in the proband. To rule out the possibility of a submicroscopic deletion, we analysed the long-range structure of the 15q11q13 region by pulsed-field gel electrophoresis (PFGE). This showed that probe IR4-3R and probe 34 detect the same large unaltered 2,500-kb *NorI* fragment in PWS2 (Fig. 3), which provides strong evidence that the absence of a paternal allele specific for probe 34 does not represent a microdeletion. Furthermore, we detected a band corresponding to a fragment of the same size (2,500 kb) in the mother, despite differential methylation leading to partial




FIG. 2 Maternal uniparental disomy in a PWS patient with apparently normal chromosomes 15. *a*, Two pairs of intact chromosomes 15 of PWS2. Previous reports have indicated these chromosomes as being intact<sup>13</sup> or deleted 15q (ref. 14); in the deletion report a polymorphic shortening of 15q11 (ref. 7) was probably detected, because we show here at a higher level of resolution that the chromosomes are intact. *b*, Maternal origin of the two chromosomes 15 in PWS2. DNA was digested with *ScaI*. PWS2 (lane 1) inherits a maternal 6.3-kb allele (lane 3) but no paternal 6.5-kb allele (lane 2) for probe 34 (D15S9). Each individual is heterozygous for IR10-1 (D15S12; 16- and 12.5-kb alleles), which is from the same 15q11q13 region as probe 34 (ref. 10). The band at 10 kb is constant, C, for probe 34 in all individuals<sup>5</sup>. *c*, PWS2 inherits maternal alleles only for probe IR39d (D15S18). DNA was digested with *SacI* and hybridized with the 15q11q13-specific probe IR39d. PWS2 (lane 1) inherits his mother's (lane 3) 14-kb alleles and not his father's (lane 2) 8.5-kb alleles. *d*, Two copies of the maternal allele for probe 34 in PWS2. *HindIII*-digested DNAs from PWS2 (lane 2), mother M2 (lane 3), father F2 (lane 4) and a PWS patient displaying both a cytological and molecular deletion of 15q11q13 (ref. 5; lane 1, HS2) were hybridized simultaneously with probes 34 and H2-26 (D13S28)<sup>31</sup>. Probe H2-26 from chromosome 13 serves as a reference for quantitation of copy number<sup>10</sup>. Probe 34 shows a reduced hybridization intensity relative to H2-26 only in HS2, which is consistent with a deletion of 15q11q13 in this individual. No reduction in probe 34 hybridization intensity is observed in PWS2 or his parents. To quantitate the intensity of the hybridization bands, the autoradiogram was scanned with an LKB Ultrascan XL laser densitometer (Pharmacia LKB Biotechnology). The area ratio of the probe 34 to probe H2-26 peaks was calculated for each lane. The number of copies of probe 34 per genome was obtained by setting the ratio in lane 3 (M2, mother) to 2 and normalizing the values in the other lanes to this value. The number of copies of probe 34 per genome determined by this method were 1.0 in lane 1 (the HS2

deletion), 1.8 in lane 2 (PWS2) and 2.2 in lane 4 (F2, father). *e*, Exclusion of non-paternity for PWS2. One maternal 5.2-kb (lane 2) and one paternal 2.9-kb (lane 3) *PvuII* allele are inherited by PWS2 (lane 1) for the 3'HVR. METHODS. DNA and cytogenetic studies were as for Fig. 1. Probe IR39d is a subclone of IR39 (ref. 10), which lacks repetitive sequences. PWS2 has been described before as patient number 5 (ref. 14) and patient number 31 (ref. 13).

*NotI* digestion in this region of the genome (Fig. 3) and in other normal individuals (results not shown). We again excluded the possibility of non-paternity using the 3'HVR fragment, which detected an entirely different set of alleles in the PWS2 family (Fig. 2e) from those that it detected in the PWS1 family (Fig. 1d).

The results presented here indicate that both PWS probands inherited two different, intact chromosome 15q11q13 regions from their mothers. Maternal heterodisomy—a newly defined form of uniparental disomy<sup>15,16</sup>—for at least the critical region of chromosome 15, could thus have an aetiological role in PWS. The extent of disomy remains to be determined by genetic analysis with other markers. The disomy extends beyond the PWS critical region identified from deletion studies because locus CMW-1, which maps distal to the deletion region, is




FIG. 3 PFGE provides evidence for the lack of a submicroscopic deletion in PWS patients with uniparental disomy. *NotI* fragments are shown for probes IR4-3R (D15S11) (left) and 34 (right) in PWS2 (lane 3), M2 (lane 2) and F2 (lane 1). Multiple bands are indicative of partial cleavage at restriction sites when CpG is methylated<sup>32</sup> and a similar pattern of multiple bands has been observed in other normal controls (results not shown). PWS2 shares a normal 2,500-kb band with his mother, although in the latter this results from partial digestion at a *NotI* site in this region of the genome. The same 2,500-kb band has been detected in other normal individuals. The larger three fragments are common to probes IR4-3R, 189-1 (results not shown) and 34, showing that these loci are closely linked in region 15q11q13. These probes and a 1,300-kb fragment detected with probes 3-21 and IR10-1 (unpublished data) are deleted in all PWS<sup>10</sup> and AS<sup>6</sup> patients with cytological deletions. Additional PFGE analysis (results not shown) also indicates that PWS2 displays PFGE restriction fragments of normal size for IR10-1 in a *NotI* digest (1,300 kb) and for IR39d in a *Bss*III digest (250 kb). PWS1 also seems to be intact for the probe 34 and 3-21 loci because the restriction fragments detected using *NotI* and *Bss*III are unaltered compared with those of normal individuals.

**METHODS.** PFGE analyses were performed as described previously<sup>33</sup> using a modification of techniques for DNA preparation and digestion in agarose blocks<sup>34</sup>. Electrophoresis was performed in 0.8% agarose in TBE (89 mM Tris buffer, pH 8.0, 89 mM boric acid, 2 mM EDTA) at 22 °C and was divided into three 48-h intervals during which the forward polarity pulse durations were varied from 75–600 s, 600–2,500 s and 2,500–3,000 s and field intensities were set at +2, +1.7 and +1.3 V cm<sup>-1</sup>, respectively. The ratios of the duration and intensity of the inverse polarity pulse relative to the forward were 0.4 and 0.5, respectively.

disomic in PWS1. Several predictions can be made about an association between uniparental disomy and the aetiology of PWS. First, there should be no chromosome deletion, as is shown in this study by using both cytogenetic and molecular genetic techniques. Additional PFGE mapping data for PWS1 and PWS2 (Fig. 3 legend) confirmed that a large part (4–5 megabases) of the 15q11q13 region is intact in these two patients. Second, there should be no specific gene mutation. Although point mutations cannot be ruled out until the gene(s) responsible for PWS are isolated, these seem unlikely because each case displays two independent maternal contributions to region 15q11q13 and the disorder is genetically dominant, which would rule out mechanisms such as the uncovering of a recessive mutation. Finally, the frequency of maternal disomy should be high in PWS patients with normal chromosomes. Consistent with this prediction, our preliminary data on four additional families strongly indicates that all of the PWS patients show maternal uniparental disomy.

It seems that the clinical phenotype of the two PWS probands arises from the absence of a paternal contribution to region 15q11q13, rather than from a specific gene mutation. This implies functional differences in alleles of a gene or genes from this region of the genome that depend on the sex of the transmitting parent (genetic imprinting<sup>18–21</sup>). Under this scheme, normal human development would require genetic input from both parents, whereas the absence of a paternal contribution to region 15q11q13, whether by paternal deletion<sup>4,5</sup> or maternal uniparental disomy (as demonstrated here), would result in PWS. Consistent with this hypothesis is our recent finding of differential transmission of parental alleles in PWS and AS (ref. 6). It is conceivable, therefore, that the absence of a maternal contribution to the same 15q11q13 region could result in a different disorder, AS.

Experimental evidence for genetic imprinting has been largely restricted to the mouse in which a requirement for a contribution of both the maternal and paternal genome for normal development has been demonstrated<sup>18–22</sup>. Transgenic studies have provided preliminary evidence that the chromatin alteration involved in the differential modification and expression of parental alleles is DNA methylation<sup>18–22</sup>, a process also associated with transcriptional regulation of gene activity<sup>23</sup>. In humans, evidence of a developmental requirement for both parental genomes is provided by the finding that only the paternal genome is present in complete hydatidiform moles<sup>24</sup>. Furthermore, differential transmission of parental alleles in several disorders such as Huntington's chorea<sup>18,25</sup>, and somatic changes in childhood tumours such as Wilms tumour, osteosarcoma (reviewed in ref. 26) and rhabdomyosarcoma<sup>27</sup> could involve genome imprinting. Growth failure in two individuals with isodisomy<sup>15</sup> for chromosome 7 (refs 16, 28) could indicate genetic imprinting<sup>16</sup>, although the aetiology of cystic fibrosis (CF) in these patients is homozygosity for a mutant CF allele inherited from the mother<sup>16,28</sup>.

In conclusion, the association of PWS with maternal uniparental disomy for region 15q11q13 implicates a role for genetic imprinting in the aetiology of the PWS phenotype. Similar phenomena could contribute to other clinical syndromes. Isolation of the gene(s) responsible for PWS and other disorders in which genetic imprinting plays a part might be possible by identifying a gene that shows a differential modification of its parental alleles. Our findings for PWS represent a step towards understanding the developmental basis of this common human genetic disorder. □

Received 31 July; accepted 13 October 1989.

- Cassidy, S. B. *Curr. Problems Pediatr.* **14**, 1–55 (1984).
- Butler, M. G. *Am. J. med. Genet.* (in the press).
- Ledbetter, D. H., Greenberg, F., Holm, V. A. & Cassidy, S. B. *Am. J. med. Genet.* **28**, 779–790 (1987).
- Butler, M. G., Meaney, F. J. & Palmer, C. G. *Am. J. med. Genet.* **23**, 793–809 (1986).
- Nicholls, R. D. et al. *Am. J. med. Genet.* **33**, 66–77 (1989).
- Knoll, J. H. M. et al. *Am. J. med. Genet.* **32**, 285–290 (1989).

7. Pembrey, M. et al. *J. Med. Genet.* **26**, 73-77 (1989).
8. Donlon, T. A. *Hum. Genet.* **80**, 322-328 (1988).
9. Donlon, T. A., Lalonde, M., Wyman, A., Bruns, G. & Latt, S. A. *Proc. natn. Acad. Sci. U.S.A.* **83**, 4408-4412 (1986).
10. Tantravahi, U. et al. *Am. J. med. Genet.* **33**, 78-87 (1989).
11. Rich, D. C., Witkowski, C. M., Summers, K. M., van Tuinen, P. & Ledbetter, D. H. *Nucleic Acids Res.* **16**, 8740 (1988).
12. Higgs, D. R. et al. *Proc. natn. Acad. Sci. U.S.A.* **83**, 5165-5169 (1986).
13. Butler, M. G. & Meaney, F. J. *Am. J. med. Genet.* **26**, 445-455 (1987).
14. Butler, M. G. *Am. J. hum. Genet.* **45**, 140-146 (1989).
15. Engel, E. *Am. J. med. Genet.* **6**, 137-143 (1980).
16. Spence, J. E. et al. *Am. J. hum. Genet.* **42**, 217-226 (1988).
17. Warburton, D. *Am. J. hum. Genet.* **42**, 215-216 (1988).
18. Solter, D. A. *Rev. Genet.* **22**, 127-146 (1988).
19. Sapienza, C. *Annals N.Y. Acad. Sci.* (in the press).
20. Monk, M. *Genes Dev.* **2**, 921-925 (1988).
21. Holliday, R. *Sci. Am.* **260**, 60-73 (1989).
22. Surani, M. A., Reik, W. & Allen, N. D. *Trends Genet.* **4**, 59-62 (1988).
23. Cedar, H. *Cell* **53**, 3-4 (1988).
24. Kajii, T. & Ohama, K. *Nature* **268**, 633-634 (1977).
25. Reik, W. *J. med. Genet.* **25**, 805-808 (1988).
26. Reik, W. & Surani, M. A. *Nature* **338**, 112-113 (1989).
27. Scrabble, H. et al. *Proc. natn. Acad. Sci. U.S.A.* (in the press).
28. Voss, R. et al. *Am. J. hum. Genet.* **45**, 373-380 (1989).
29. Yunis, J. J. *Science* **191**, 1268-1270 (1976).
30. Seabright, M. *Lancet* **ii**, 971-972 (1971).
31. Lalonde, M. et al. *Cancer Genet. Cytogenet.* **13**, 283-295 (1984).
32. Fischel-Ghodsian, N., Nicholls, R. D. & Higgs, D. R. *Nucleic Acids Res.* **15**, 6197-6207 (1987).
33. Lalonde, M., Noolandi, J., Turmel, C., Rousseau, J. & Slater, G. W. *Proc. natn. Acad. Sci. U.S.A.* **84**, 8011-8015 (1987).
34. Schwartz, D. C. & Cantor, C. R. *Cell* **37**, 67-75 (1984).

ACKNOWLEDGEMENTS. We are grateful to Dr L. Kunkel for his critical reading of the manuscript, D. Shook and K. Glatt for technical assistance, B. Woolf for help in preparation of the manuscript, Dr D. Ledbetter for providing the CMW-1 probe and Dr C. Sapienza for communicating results and hypotheses before their publication. We also thank Dr S. Kapur for clinical evaluation of one of the patients. This work was supported in part by the NIH. M.L. is a senior associate of the Howard Hughes Medical Institute.

## Requirement for integrins during *Drosophila* wing development

Danny L. Brower & Sharon M. Jaffe

Department of Molecular and Cellular Biology, Biosciences West Building, University of Arizona, Tucson, Arizona 85721, USA

THE position-specific (PS) integrins of *Drosophila*<sup>1,2</sup> are highly homologous to vertebrate integrins<sup>3-5</sup>, most of which are cell-surface receptors for extracellular matrix components<sup>6,7</sup>. Integrins are heterodimers, each consisting of noncovalently associated  $\alpha$ - and  $\beta$ -subunits. As for the subfamilies of vertebrate integrins, the same  $\beta$ -subunit is found in both *Drosophila* PS integrins, combined with a specific  $\alpha$ -subunit to generate either a complete functional PS1 or PS2 integrin<sup>1,5,8</sup>. Both  $\alpha$ - and  $\beta$ -subunits are large transmembrane proteins (relative molecular masses >100,000). Either one or both of these two PS integrins are expressed in most fly tissues during development. A particularly intriguing pattern of expression is found in the mature wing imaginal disc, where the PS1 integrin is expressed primarily on the presumptive dorsal wing epithelium, and the PS2 integrin is found almost exclusively on the ventral epithelium<sup>1</sup>. Immediately after pupariation, the central wing pouch evaginates, folding along its centre to appose the epithelia that will secrete the dorsal and ventral surfaces of the adult wing blade<sup>9</sup>. Here we report the results of a genetic analysis indicating that both of the PS integrins are required to maintain the close apposition of the dorsal and ventral wing epithelia during morphogenesis. Also, we conclude that the integrins are not necessary for the maintenance of the cell lineage restriction between the two presumptive wing surfaces in the developing imaginal disc<sup>10-12</sup>.

The  $\alpha$ -subunit of the PS2 integrin seems to be encoded by the *inflated* (*if*) locus on the X chromosome<sup>13</sup>, and null mutations at *if* cause embryonic lethality (M. Wilcox, A. DiAntonio and M. Leptin, manuscript submitted; see also Fig. 2 legend). We examined *if*<sup>3</sup> mutant larval tissues by using monoclonal antibodies directed against the PS integrins (anti-PS antibodies), and our results suggest that the *if*<sup>3</sup> allele is a regulatory mutation

at the locus. Homozygous *if*<sup>3</sup> late third-instar larvae (just before pupariation and disc evagination) had apparently normal levels of PS2 integrin on most tissues (for example, muscles, salivary glands), but displayed greatly reduced levels of integrin on the surfaces of some imaginal disc cells (Fig. 1*b*). Specifically, there was relatively little PS2 integrin  $\alpha$ -subunit in the ventral region of the wing pouch; these cells normally express the protein at very high levels. PS2 integrin expression was not so severely reduced in other regions of the disc, such as the peripodial membrane. The expression of the PS1 integrin in the wing disc seemed to be unaffected by *if*<sup>3</sup> (not shown).

Although all of the mutant wing discs examined were clearly affected, there was variability in the extent of PS2 integrin reduction (for example, Fig. 1*b, c*). In the wing pouch, mutant discs typically stained at higher levels anteriorly, and often displayed significant levels of staining with anti-PS2 antibody dorsally (Fig. 1*c*). Similar antero-posterior asymmetry and dorsal staining are characteristic of wild-type PS2 integrin expression in mid-third instar discs<sup>14</sup>, about 24 h before pupariation. It is interesting that the pattern of PS2 integrin expression in *if*<sup>3</sup> discs seemed relatively normal at the mid-third larval instar (not shown), and it is possible that this allele simply does not allow the cells of the pouch to progress to the more mature pattern of expression.

Adult flies bearing the *if*<sup>3</sup> allele display wing blisters, in which the dorsal and ventral surfaces of the wing blade are separated<sup>13</sup>. We found that the penetrance and expressivity of the *if*<sup>3</sup> blisters were variable (Fig. 2); this is not surprising in light of the variability of the immunofluorescence results. Nonetheless, these data indicate that the PS2 integrin is important for maintaining the close apposition of the dorsal and ventral epithelia during morphogenesis of the adult wing.

That this is the case was confirmed by clonal analysis studies<sup>15</sup> of flies with mutations in the *mysospheroid* gene (*mys*). The *mys* locus encodes the common PS integrin  $\beta$ -subunit<sup>4,5</sup>, so null mutations at this locus would be expected to eliminate both PS1 and PS2 integrins<sup>5</sup>. Flies homozygous for null mutations of *mys* die as embryos (see refs 16 and 17 for descriptions of the *mys* lethal phenotype), so to examine the role of integrins in wing morphogenesis, we made clones of cells homozygous for a null mutation of *mys*, along with the flanking markers *yellow* (*y*) and



FIG. 1 Anti-PS2-antibody immunofluorescence of the basal surfaces of late third-instar wing imaginal discs. At this stage, the disc consists primarily of a single layered but highly folded columnar epithelium of ~50,000 cells<sup>22</sup>. All micrographs are centred on the 'pouch' region, which evaginates into a flat sac before secreting the cuticle of the adult wing blade<sup>9</sup>. Images are reverse-contrast, so staining is dark. *a*, Wild type. PS2 integrin is found at high concentration throughout the ventral (upper) region of the wing pouch. The sharp horizontal boundary of staining across the pouch marks the line along which the epithelium will fold during evagination (that is, the presumptive wing margin). *b*, Mutant for *if*<sup>3</sup>. PS2 expression is greatly reduced in the pouch. Relatively high levels of staining remain in more peripheral areas, especially along the anterior edge of the pouch (arrow). *c*, Mutant for *if*<sup>3</sup>. Higher magnification of another disc with a less extreme phenotype, showing residual PS2 in the pouch. Typically, expression is greater in the anterior (left) half of the pouch, and is often seen dorsally (arrow) as well as ventrally. Scale bar, 50  $\mu$ m.

METHODS. Mutants for *if*<sup>3</sup> were grown as a homozygous stock at 25 °C. Immunofluorescence of wing discs was performed with an anti-PS2 (CF.2C7) monoclonal antibody and fluorescein-conjugated goat anti-mouse antibody (Antibodies Inc.) as previously described<sup>1</sup>. Images were detected with an ISIT video camera, and stored on video tape.